


Laminar burning velocities



References

- [1] F. Egolfopoulos, C. Law, An experimental and computational study of the burning rates of ultra-lean to moderately-rich H₂/O₂/N₂ laminar flames with pressure variations, Symp. (Int.) Comb., 23 (1990) 333–340.
- [2] D. R. Dowdy, D. B. Smith, S. C. Taylor, A. Williams, The use of expanding spherical flames to determine burning velocities and stretch effects in hydrogen/air mixtures, Symp. (Int.) Comb., 23 (1990) 325 – 332.
- [3] S. D. Tse, D. L. Zhu, C. K. Law, Morphology and burning rates of expanding spherical flames in H₂/O₂/inert mixtures up to 60 atmospheres, Proc. Comb. Inst, 28 (2000) 1793 – 1800.
- [4] O. C. Kwon, G. M. Faeth, Flame/stretch interactions of premixed hydrogen-fueled flames: measurements and predictions, Comb. Flame, 124 (2001) 590 – 610.
- [5] N. Lamoureux, N. Djebaili-Chaumeix, C.-E. Paillard, Laminar flame velocity determination for H₂-air-He-CO₂ mixtures using the spherical bomb method, Exp. Therm. Fluid Sci., 27 (2003) 385–393.
- [6] S. Verhelst, R. Woolley, M. Lawes, R. Sierens, Laminar and unstable burning velocities and Markstein lengths of hydrogen-air mixtures at engine-like conditions, Proc. Comb. Inst., 30 (2005) 209 – 216.